



## RESPONSE OF SQUASH PLANTS GROWN UNDER SANDY SOIL AND HIGH IRRIGATION WATER SALINITY CONDITIONS TO FOLIAR SPRAY WITH POTASSIUM SILICATE AND INOCULATION WITH MYCORRHIZA

Alaa El-Den H. Roshdy<sup>1,\*</sup> and Noura M. Taha<sup>2</sup>

<sup>1</sup>Horticultur Department, Faculty of Agriculture, Damanhour University, **Egypt.**

<sup>2</sup>Horticultur Department, Faculty of Agriculture, Ain Shams University, **Egypt.**

\*Corresponding author: [alaa.roshdy@agr.dmu.edu.eg](mailto:alaa.roshdy@agr.dmu.edu.eg) Received: 10 Aug. 2020; Accepted: 12 Sep. 2020

**ABSTRACT:** An experiment was carried out at Wadi El-Natroon district Behiera Governorate, during the growth summer seasons of 2019 and 2020 to investigate the effect of inoculation with mycorrhiza (M) and/or foliar application of silicon as potassium silicate (KS) to improve squash plants growth parameters and yield and its components as well as fruits nutrient quality, which confronting the saline conditions in new reclaimed soils. The irrigation of squash plants was with well water that has salinity ranged between 3.4–3.6 dSm<sup>-1</sup>. The results revealed decreasing of all growth, yield and fruits nutrients quality characters as shown in control (well water). However, the application of mycorrhiza inoculum in squash seeds accompanied with foliar application of KS at 4 mL<sup>-1</sup> every 10 days after transplanting in the open field could be an effective procedure for significant enhancing the performances of squash plants grown under similar unfavorable conditions of increasing irrigation water salinity.

**Key words:** squash, *Cucurbita pepo*, salinity, mycorrhiza, potassium silicate, growth, yield.

### INTRODUCTION

Due to urban transgression, the amount of traditional agricultural land in the Nile Delta has declined sharply in recent decades. In particular, in the last decade, Egypt has witnessed a steady increase in the agricultural area, which was by the reclamation of new land areas that irrigated by groundwater in most cases (Quosy, 2019; Radwan, 2019). However, it is well known that the dominant problem on such new reclaimed areas in Egypt is the salinity (Chhabra, 2017), which narrow the chances of growing many kinds of cash valuable vegetable crops, which exhibit significant reduction in productivity, especially if irrigated with saline water (Machado and Serralheiro, 2107).

Salinity stress effects could be devoted in three aspects e.g., osmotic stress, specific stress and oxidative stress; however, the oxidative stress is considered to be the most destructive (Tanveer and Shabala, 2018). High salinity not only declines plant growth, biomass, yield, photosynthesis, and water use efficiency, but also resulted in

physiological drought, ion toxicity in plants and imbalance nutrient uptake, which, consequently, reducing the agricultural productivity and yields of the stressed crop plants (Safdar, 2019; Shahid, 2018 and Wungrampha, 2018).

Among the domesticated Cucurbits, the squash (*Cucurbita pepo* L.) seeds was archeologically discovered in Mexico that are more than 9,000 years old (Decker, 1988). According to the salinity tolerance scale, Squash is categorized as a moderately tolerant with thresholds estimated by 4.7 and 3.1 dSm<sup>-1</sup> for soil and water salinity, respectively (Ayers and Westcot, 1985).

There are many procedures were examined by researchers that could be effective in order to ameliorate the dangerous effects of salinity stress, including the inoculation with a beneficial microorganism, such as mycorrhiza fungi or treatment with some nutrients, such as silicon (Machado and Serralheiro, 2107; Plaut *et al.*, 2013).

Inoculation with mycorrhiza is considered as an effective strategy in alleviating plant growth under saline conditions. Many of investigations have exhibited that mycorrhiza inoculations improve growth by diminishing the development of reactive oxygen species (ROS) and alleviating antioxidant defense enzymes activities under salt stress conditions (Wu, 2014). Accordingly, mycorrhiza inoculation was found to increase the different physiological parameters, e.g. chlorophyll content, photosynthetic process and nutrient uptake, especially N, P and K with association with decreasing Na uptake (Begum *et al.*, 2019; Rahman *et al.*, 2019 and Ramadan, 2019), which reflects on enhancing growth and yield of plants grown under salinity stress.

Also, the exogenous application of silicon (Si) has been a recent eco-friendly approach to enhance the salinity stress response in plants (Almeida, 2017). The application of silicon on the salinity stressed plants was found to ameliorate the deleterious effect of salinity stress in many ways as decreasing the rate of Na uptake, increasing root water uptake, reinforcement the cell walls, improve the efficiency of osmotic adjustment and enhancing the photosynthesis (Zhu, 2019). The investigations about the role of Si in vegetable dicots, however, are still far behind cereals (Kaushik and Saini, 2019).

Among the silicon-based fertilizers, potassium silicate ( $K_2O_3Si$ ) was the most popular water-soluble silicate fertilizer that that used as foliar spray (Laane, 2018), which provides silicon (25%  $SiO_2$ ) to the treated plants in addition to few amounts of potassium (10%  $K_2O$ ). The use of this soluble silicate fertilizer as foliar application is due to their high cost for soil application (Yan *et al.*, 2018).

Moreover, there were many research papers demonstrate a synergistic effect between silicon and mycorrhiza, i.e. the mycorrhiza formation and functions were increased by Si, and; in turn, Si uptake was increased due to mycorrhization (Frew, 2017; Moradtalab, 2019); which increase their effectiveness in alleviating the consequences of abiotic stress of the treated plants e.g., drought (Azam Eidi and Najafabadi, 2014; Hajiboland *et al.*, 2018), heavy metals (Garg and Singh, 2018) and salinity (Garg and Bhandari, 2016) as well as under normal conditions (Hajiboland *et al.*, 2018).

The aim of this study was to investigate the possibility of inoculation with mycorrhiza (M) and/or foliar application of silicon as potassium silicate (KS) to enhance the ability of squash plants

to confronting the saline conditions in new reclaimed soils.

## MATERIALS AND METHODS

The field experiment was conducted in the summer seasons of 2019 and 2020 in a private farm, at Wadi El-Natroon region, Behiera Governorate, Egypt, under open field conditions in a sandy soil where this farm is suffering from salinity of irrigation water more than soil as shown in Tables 1 and 2.

Tables (1 and 2) showed the different physical and chemical parameters of the soil and irrigation water in the experimental location. The analyses were determined according to Dewis and Freitas (1970).

The cultivar used for the experiment was "Aziad" squash hybrid, which was sown in transplant trays for 25 days, then seedling transplanted in the experimental field at 18<sup>th</sup> and 22<sup>th</sup> February of 2019 and 2020, respectively.

This experiment included six treatment. The experimental design was split plots with three replicates. The mycorrhiza inoculation treatments (M<sub>-</sub> and M<sub>+</sub>) occupied the main plots and the potassium silicate treatments (KS<sub>0</sub>, KS<sub>1</sub> and KS<sub>2</sub>) occupied the sub-plots at each replicate. Each subplot consisted of four rows, 6 m long and 70 cm width for each; also, each plot was surrounded with guard row, 70 cm width to minimizing the interferences between treatments.

The treatments of mycorrhiza inoculum (*Glomus* spp.) were applied as a suspension of spores and root pieces after five days from sowing squash seeds in transplant trays. The mycorrhiza inoculum was derived from faculty of Agriculture, Saba Basha, Alex. Univ., Biofertilization Unit; whereas, the silicon treatments were applied as potassium silicate ( $K_2O_3Si$ ). Three graded levels of potassium silicate (KS) were used as foliar application, e.g. control (well water), 2 and 4 mL<sup>-1</sup> that denoted as KS<sub>0</sub>, KS<sub>1</sub> and KS<sub>2</sub>, respectively. The foliar application of each examined concentration was applied after ten days from transplanting and was repeated every ten days until 15 days before the end of the season. All the agricultural practices that recommended for the squash production in the new reclaimed soils under drip irrigation system, e.g. fertilization, pest control and irrigation schedules, were followed.

**Table 1. Physical and chemical analysis of the experimental soil in 2019/2020 seasons of study**

Physical analysis*	2019	2020
Sand	93	92
Clay	4	5
Silt	3	3
Soil texture	Sand	Sand
Chemical analysis*		
pH	8.4	8.5
EC (dSm <sup>-1</sup> )	3.5	3.7
CaCO <sub>3</sub> (%)	5.4	5.6
Macro elements (ppm)		
N	16	14
P	17	20
K	61	63
Cations (meq L <sup>-1</sup> )		
Ca <sup>+2</sup>	3.1	3.3
Mg <sup>+2</sup>	2.4	2.7
K <sup>+</sup>	1.6	1.6
Na <sup>+</sup>	3.6	3.9
Anions (meq L <sup>-1</sup> )		
Cl <sup>-</sup>	2.4	2.8
SO <sub>4</sub> <sup>-</sup>	7.5	7.6
CO <sub>3</sub> <sup>-</sup>	--	--
HCO <sub>3</sub> <sup>-</sup>	0.8	1.1
<b>SAR</b>	2.2	2.3

\* These analyses were carried out at the Soil Reclamation and Agric. Engineering Dept., Agric. Fac., Damanhour University.

**Table 2. Chemical analysis of the irrigation water (well water) used in the experiment in 2019 and 2020 seasons**

Chemical analysis*	2019	2020
pH	7.5	7.7
EC (dSm <sup>-1</sup> )	3.5	3.7
Cations (meq L <sup>-1</sup> )		
Ca <sup>+2</sup>	3.0	3.2
Mg <sup>+2</sup>	3.3	3.3
K <sup>+</sup>	2.7	3.1
Na <sup>+</sup>	19.8	21.4
Anions (meq L <sup>-1</sup> )		
Cl <sup>-</sup>	17.8	19.4
SO <sub>4</sub> <sup>-</sup>	6.2	6.5
CO <sub>3</sub> <sup>-</sup>	--	--
HCO <sub>3</sub> <sup>-</sup>	4.8	5.1
<b>SAR</b>	11.2	11.9

\* These analyses were carried out at the Soil Reclamation and Agric. Engineering Dept., Agric. Fac., Damanhour University.

The measured characters were divided to four groups as follow:

#### Vegetative growth characters

After 40 days from transplanting, stem length, leaves number plant<sup>-1</sup>, fresh and dry weight of shoot (stems and leaves) were recorded.

#### Leaf chlorophyll and mineral constitute

Total Chlorophyll: was measured after 40 days from transplanting by non-destructive method using SPAD-502 chlorophyll meter (Markwell *et al.*, 1995).

Leaves mineral contents: minerals were determined as illustrated by Kalra (1998); nitrogen (N) was determined using micro-kjeldahl method, phosphorus (P), potassium (K) and sodium (Na) were estimated using flam photometer. Moreover, K<sup>+</sup>/Na<sup>+</sup> ratio that was calculated.

#### Yield and its components

After 40 days from transplanting, the fruits were harvested from each plot every 3 days and the following data were recorded: Fruit number plant<sup>-1</sup>, average fruit weight (which was derived from the average of ten fruits weight in grams), fruit yield plant<sup>-1</sup> and yield faddan<sup>-1</sup>.

#### Fruit quality

The macro- and micro-elements of N, P, K, Ca, Cu, Zn and Fe in squash fruits were determined following the steps that demonstrated by Kalra (1998).

#### Statistical analysis

All recorded data were statistically analyzed by CoStat software (COSTAT, 2005) version 6.4 from cohort software. Least significant difference test (LSD) was applied at 0.05 confidence level to compare means of different treatments using the same program.

## RESULTS AND DISCUSSION

#### Vegetative growth characters

Data in Table 3 showed that the M inoculation provide significant increases in all vegetative growth characters (stem height, leaves number plant<sup>-1</sup>, shoot fresh weight and shoot dry weight) if compared with uninoculated plants irrigated with saline water without inoculation.

Concerning the effect of KS treatments, the results revealed significant enhancing effect for the KS foliar treatments on vegetative growth traits of squash plants grown under saline conditions when compared with control (KS<sub>0</sub>), in both seasons of study. The foliar application of KS<sub>2</sub> was the best stem height, leaf number, shoot fresh and dry weight.

The interaction between M and KS indicated that the foliar application with KS<sub>2</sub> of the squash plants inoculated with M was the superior application over all the other interaction combinations, in both seasons of study.

These results revealed significant ameliorative effect for each of mycorrhiza inoculation or potassium silicate foliar application on vegetative growth of squash plants grown under saline conditions, in both seasons of study. At the same trend of the results of this study, many other researchers illustrated the important effect of mycorrhiza inoculation on improving the vegetative growth characters of plants grown under saline conditions (Cantrell and Linderman, 2001 on lettuce and onion; Colla *et al.*, 2008 on zucchini and Haghghi and Barzegar, 2017 on sweet pepper).

The ameliorative effect of M inoculation could be due to occurring an enhancement of activity of different antioxidant enzymes in salt affected plants that indicating lower oxidative damage (Abdel Latif *et al.*, 2011). Also, the M fungi were found to produce different types of hormone-like substances e.g., gibberellins and cytokinins (Amballa and Bhumi, 2017). In addition, mycorrhiza has the ability to extending the absorbing zones of host plant roots that ensure a reduction in nutrient loss (Cavagnaro, 2015). Consequently, the mycorrhiza has the ability to enhance all of vegetative growth characters of the host plants.

Concerning the KS effect, it could due to the effect e.g., the effect of potassium and the effect of silicon. potassium is a vital for plant survival under physiological and stress conditions because its role in many physiological and biochemical processes (Wang *et al.*, 2013). Moreover, Shafeek *et al.* (2013) on cucumber and Fekry (2016) on summer squash reported that potassium foliar application enhanced the vegetative growth characters. On the other side, many researchers demonstrated the important role of silicon on alleviating the growth parameters of many crop plants under normal or stressed conditions by enhancing water relations, decreasing the Na uptake, increasing photosynthesis rate, nutrients uptake and antioxidant enzymes (Laane, 2018) as well as impacting the phytohormones that commonly involved in controlling stress conditions (Luyckx *et al.*, 2017). Moreover, many research studies revealed that the spraying of potassium silicate has favorable effect

on growth parameters of many of vegetable crops (Amirossadat *et al.*, 2012 on cucumber; Hussein and Muhammed, 2017 on eggplant; Merwad, 2018 on pea; Abd-Elaziz *et al.*, 2019 and Shehata and Abdelgawad, 2019 on squash and Yaghubi *et al.*, 2019 on strawberry).

The interaction between M inoculation and KS application was found to give more ameliorative effect on vegetative growth characters of squash plants than the single treatments, which illustrated by the increment percentages in vegetative growth characters comparing to control in the case of single or combined treatments, in both seasons of study. These enhancement effect of the combined treatment could be a result of some kind of synergistic relationship between M and KS as suggested by Frew *et al.* (2017); and Moradtalab *et al.* (2019). In addition, Garg and Bhandari (2016) found in *Cicer arietinum* genotypes that the mycorrhiza increased the endogenous amounts of silicon and the application of silicon enhanced the efficiency of mycorrhiza fungi. Moreover, they suggested that the Si-mycorrhiza application could be effective treatment for overcoming the salinity conditions.

#### Leaf chlorophyll and mineral constitute

As shown in Table 4, the results revealed significant effect of the M inoculation on leaf chlorophyll as well as N, P, K and K/Na ratio but decreased Na, in both seasons of study. The increased values of the uninoculated squash plants were 23.69% for chlorophyll (SPAD), 59.88% for N, 39.25% for P, 59.17% for K and 105.15% for K/Na. However, the leaves content of Na was decreased due to inoculation with M, in both seasons of study with percentage of -23.15% compared with control as both seasons average.

According the mean values of the KS foliar applications, with the exception of Na, the KS<sub>2</sub> treatment gave the highest mean values of each of chlorophyll, N, P, K and K/Na ratio in squash leaves, in both seasons of study. However, the KS<sub>2</sub> treatment decreased, significantly, the squash leaf values of Na, in both seasons of study.

**Table 3. Effect of mycorrhiza inoculation, potassium silicate foliar applications and their interaction on vegetative growth of squash plants 2019 and 2020 seasons**

Treatments	2019*			2020*		
	M <sub>-</sub>	M <sub>+</sub>	Mean	M <sub>-</sub>	M <sub>+</sub>	Mean
Stem length (cm)						
KS <sub>0</sub>	12.72 e	19.78 b	<b>16.25 C</b>	13.16 f	17.58 c	<b>15.37 C</b>
KS <sub>1</sub>	16.27 d	20.70 b	<b>18.49 B</b>	15.95 e	19.05 b	<b>17.50 B</b>
KS <sub>2</sub>	18.34 c	23.21 a	<b>20.77 A</b>	16.75 d	21.59 a	<b>19.16 A</b>
<b>Mean</b>	<b>15.78 B</b>	<b>21.23 A</b>		<b>15.29 B</b>	<b>19.41 A</b>	
Leaf number plant <sup>-1</sup>						
KS <sub>0</sub>	9.36 f	14.97 c	<b>12.17 C</b>	8.89 f	13.89 c	<b>11.39 C</b>
KS <sub>1</sub>	11.41 e	15.97 b	<b>13.68 B</b>	10.25 e	15.33 b	<b>12.79 B</b>
KS <sub>2</sub>	13.30 d	17.99 a	<b>15.65 A</b>	12.22 d	16.89 a	<b>14.56 A</b>
<b>Mean</b>	<b>11.36 B</b>	<b>16.31 A</b>		<b>10.45 B</b>	<b>15.37 A</b>	
Shoot fresh weight (gm)						
KS <sub>0</sub>	327.26 e	432.59 c	<b>379.92 C</b>	318.77 f	407.31 c	<b>363.05 C</b>
KS <sub>1</sub>	362.91 d	458.30 b	<b>410.60 B</b>	340.69 e	451.63 b	<b>396.16 B</b>
KS <sub>2</sub>	366.70 d	486.57 a	<b>426.63 A</b>	362.61 d	472.94 a	<b>417.78 A</b>
<b>Mean</b>	<b>352.29 B</b>	<b>459.15 A</b>		<b>340.69 B</b>	<b>443.96 A</b>	
Shoot dry weight (gm)						
KS <sub>0</sub>	46.66 d	64.62 c	<b>55.64 C</b>	45.80 e	53.93 c	<b>49.87 C</b>
KS <sub>1</sub>	58.92 c	71.66 b	<b>65.29 B</b>	47.72 d	55.76 b	<b>51.74 B</b>
KS <sub>2</sub>	60.19 c	80.81 a	<b>70.50 A</b>	49.08 d	58.46 a	<b>54.77 A</b>
<b>Mean</b>	<b>55.26 B</b>	<b>72.36 A</b>		<b>47.54 B</b>	<b>56.05 A</b>	

\* Mean values with the same alphabetical letter (s) did not differ significantly at  $p < 0.05$  confidence level according to LSD test.

Moreover, the effect of interaction between M inoculation and KS treatments was found to be significant for all characters, in both seasons of study. Also, the interaction between M<sub>+</sub> with KS<sub>2</sub> foliar application gave, the highest mean values of N, P, K and K/Na ratio, in both seasons of study, but decreased the values of Na, in both seasons of study.

The outcomes of this investigation showed that the application of mycorrhiza or potassium silicate effectively increased leaf chlorophyll content and the activity of nutrients uptake of N, P and K in parallel with decreasing the Na contents of squash leaves that was reflected on increasing the K/Na ratio comparing with the squash plants grown under saline conditions. Also, there was superiority of mycorrhiza inoculation over potassium silicate in increasing chlorophyll content and nutrients content and K/Na ratio and decreasing Na content of squash leaves.

This could be due to that the mycorrhiza provide more surface area for nutrients absorption of the host plant roots, which promotes nutrients uptake not only phosphorus but nitrogen and potassium too, comparing with uninoculated plants (Begum *et al.*, 2019). Also, the decreasing of Na content due to mycorrhiza inoculation could be due to its ability to excluding Na<sup>+</sup> ions, through discriminating its uptake from the soil or during its transfer to plants,

thus acting as a first barrier for ion selection. Consequently, the K/Na ratio as well as with the other nutrients were increased (Hammer *et al.*, 2011; Garg and Bhandari, 2016). Moreover, mycorrhiza was found to have a vital role in increasing the leaf content of photosynthetic pigments of the host plants as shown with Cantrell and Linderman (2001) on lettuce and onion and Colla *et al.* (2008) on zucchini.

Concerning the positive effect of KS, it could be due to the role of each of potassium and silicon in enhancing the ability of plant to confront salinity. Potassium plays important role in maintenance of plant grown under different types of abiotic stresses (Römheld and Kirkby, 2010; Wang *et al.*, 2013). Also, the silicon proportion significantly affected the different nutrients uptake under normal or saline conditions as illustrated by Korkmaz *et al.* (2018) on tomato and Merwad (2018) on pea.

In addition, Abd-Elaziz *et al.* (2019) and Shehata and Abdelgawad (2019) found that the squash leaf values of N, P and K as well as chlorophyll were increased, significantly, due to the KS foliar application. Under salinity, potassium silicate was found to have ameliorating effect where Si blocks Na entrance and enhances K uptake as well as other nutrients, and significantly increasing K/Na ratio as proposed by Cantrell and Linderman

(2001) on lettuce and onion, Savvas *et al.* (2009) on zucchini and Yaghubi *et al.* (2019) on strawberry.

These may suggested that the application of KS increase the capability of mycorrhiza in acquisition

of more nutrients from the soil comparing with the only inoculated plants, which improve the endogenous nutrients profile of plants grown under salinity stress (Garg and Bhandari, 2016).

**Table 4. Effect of mycorrhiza inoculation, potassium silicate foliar applications and their interaction on leaf content of chlorophyll, N, P, K, Na and K/Na ratio 2019 and 2020 seasons**

Treatments	2019*			2020*		
	M <sub>-</sub>	M <sub>+</sub>	Mean	M <sub>-</sub>	M <sub>+</sub>	Mean
Chlorophyll (SPAD)						
KS <sub>0</sub>	34.02 d	44.98b	<b>39.50 C</b>	34.67 e	44.11 b	<b>39.39 C</b>
KS <sub>1</sub>	38.87 c	47.42 ab	<b>43.15 B</b>	36.78 d	45.74 b	<b>41.26 B</b>
KS <sub>2</sub>	41.30 c	50.83 a	<b>46.06 A</b>	41.22 c	47.55 a	<b>44.39 A</b>
Mean	<b>38.06 B</b>	<b>47.74 A</b>		<b>37.56 B</b>	<b>45.80 A</b>	
N%						
KS <sub>0</sub>	1.132 e	2.070 b	<b>1.601 C</b>	1.062 d	1.797 b	<b>1.429 C</b>
KS <sub>1</sub>	1.293 d	2.366 a	<b>1.830 B</b>	1.422 c	2.001 a	<b>1.712 B</b>
KS <sub>2</sub>	1.578 c	2.449 a	<b>2.014 A</b>	1.536 c	2.145 a	<b>1.841 A</b>
Mean	<b>1.335 B</b>	<b>2.295 A</b>		<b>1.340 B</b>	<b>1.981 A</b>	
P%						
KS <sub>0</sub>	0.270 f	0.386 c	<b>0.328 C</b>	0.263 d	0.335 c	<b>0.299 C</b>
KS <sub>1</sub>	0.297 e	0.421 b	<b>0.359 B</b>	0.275 d	0.358 b	<b>0.319 B</b>
KS <sub>2</sub>	0.316 d	0.456 a	<b>0.386 A</b>	0.279 d	0.410 a	<b>0.342 A</b>
Mean	<b>0.294 B</b>	<b>0.421 A</b>		<b>0.272 B</b>	<b>0.368 A</b>	
K%						
KS <sub>0</sub>	1.199 f	2.222 c	<b>1.710 C</b>	1.211 e	1.810 c	<b>1.511 C</b>
KS <sub>1</sub>	1.470 e	2.468 b	<b>1.969 B</b>	1.478 d	2.147 b	<b>1.812 B</b>
KS <sub>2</sub>	1.819 d	2.783 a	<b>2.301 A</b>	1.601 d	2.556 a	<b>2.079 A</b>
Mean	<b>1.496 B</b>	<b>2.491 A</b>		<b>1.430 B</b>	<b>2.171 A</b>	
Na						
KS <sub>0</sub>	0.697 a	0.533 cd	<b>0.615 A</b>	0.743 a	0.534 d	<b>0.639 A</b>
KS <sub>1</sub>	0.618 b	0.486 de	<b>0.552 B</b>	0.666 b	0.496 de	<b>0.581 B</b>
KS <sub>2</sub>	0.546 c	0.450 e	<b>0.498 C</b>	0.591 c	0.469 e	<b>0.530 C</b>
Mean	<b>0.621 A</b>	<b>0.490 B</b>		<b>0.667 A</b>	<b>0.499 B</b>	
K/Na ratio						
KS <sub>0</sub>	1.721 f	4.173 c	<b>2.947 C</b>	1.630 f	3.396 c	<b>2.513 C</b>
KS <sub>1</sub>	2.346 e	5.086 b	<b>3.716 B</b>	2.223 e	4.335 b	<b>3.279 B</b>
KS <sub>2</sub>	3.327 d	6.222 a	<b>4.775 A</b>	2.708 d	5.454 a	<b>4.081 A</b>
Mean	<b>2.465 B</b>	<b>5.160 A</b>		<b>2.187 B</b>	<b>4.395 A</b>	

\* Mean values with the same alphabetical letter (s) did not differ significantly at  $p < 0.05$  confidence level according to LSD test.

### Yield and its components

The mycorrhizal inoculation resulted in increasing percentages estimated of yield and its components by 25.33% for fruit numberplant<sup>-1</sup>, 11.89% for average fruit weight, 31.27% for fruit yield plant<sup>-1</sup> and 35.08% for fruit yield faddan<sup>-1</sup>, as average of both seasons of study if compared with uninoculated squash plants.

Also, the foliar application of KS, the treatment of KS<sub>2</sub> was the most effective concentration for

increasing all yield and its components traits of squash plants if compared with KS<sub>1</sub> or KS<sub>0</sub>, in both seasons of study. The estimated average of increasing due to the foliar application with KS<sub>2</sub> in both seasons was 17.61% for fruit numberplant<sup>-1</sup>, .825% for average fruit weight, 23.67% for fruit yield plant<sup>-1</sup> and 26.80% for fruit yield faddan<sup>-1</sup> comparing with control (KS<sub>0</sub>).

The mean values concerning the interaction between M and KS was clarified significant effect

on all yield traits, in both seasons of study. The inoculated squash plants with M and treated with KS<sub>1</sub> or KS<sub>2</sub> gave the highest interaction mean values without significant differences between them for all the yield characters except with fruit yield plant<sup>-1</sup> and fruit yield faddan<sup>-1</sup>, only in the second season where the M<sub>+</sub>+KS<sub>2</sub> treatment gave the higher mean values comparing with M<sub>+</sub>+KS<sub>1</sub>. If the relationship between M<sub>+</sub>+KS<sub>2</sub> and control, e.g. M<sub>-</sub>+KS<sub>0</sub> is taken into consideration. The average of increments in interaction mean values of M<sub>+</sub>+KS<sub>2</sub> over the control were 20.25, 22.1, 36.96 and 75.86% for fruits No. plant<sup>-1</sup>, average fruit weight, fruit yield plant<sup>-1</sup> and fruit yield faddan<sup>-1</sup>, respectively.

The enhancing of yield and its components is always considered as consequence for improvement in nutrient contents and vegetative growth parameters. From the above-mentioned results, the same trend of effect of M or KS was found on the studied yield and its components characters. Concerning the mycorrhizal effect, many researchers demonstrated the important role of such fungus in increasing the uptake of many nutrients including nitrogen, potassium, phosphorus, photosynthetic pigments as well as decreasing the uptake of sodium, which reflects on enhancing vegetative growth performance, which led to improve the yield and its component characters of the host plants especially under saline conditions (**Rahman et al., 2019 and Ramadan, 2019**).

Concerning the KS treatments, there ameliorative effect was found for yield and its components traits of squash plants. Potassium element is well known for enhancing the yield and its components of many of vegetable crops under normal (**Shafeek et al.** on cucumber, **2013**; **Fekry** on summer squash, **2016**) or under salinity stress conditions (**Römhald and Kirkby, 2010**; **Wang et al., 2013**). Also, silicon demonstrated by many researchers as a beneficial element, which was found to contribute to increasing yield and its components especially under saline conditions (**Korkmaz et al.** on tomato, **2018**). The Si effect could be because of its role in significant increase in the activities of antioxidants, decrease in the contents of electrolytic leakage percentage and adjustment of osmotic stress (**Zhu et al., 2019**), which reflects positively on

yield. Many other researchers reported the significant effect for KS foliar application on enhancing the yield and its components of the treated plants (**Hussein and Muhammed, 2017** on eggplant; **Abd-Elaziz et al., 2019** and **Shehata and Abdelgawad, 2019** on squash).

There was synergistic effect for the interaction between mycorrhiza and potassium silicate where increased the mean values of yield and its components characters as fruit No. plant<sup>-1</sup>, average fruit weight, fruit yield plant<sup>-1</sup> and fruit yield faddan<sup>-1</sup> comparing with the solo treatments. This enhancing effect of KS and AM combined applications might to be related to improving vegetative growth (Table 3) and nutrient status (Table 4), increasing the expression of the antioxidant enzymes and elevating water uptake and its use efficiency (**Garg and Bhandari, 2016**; **Moradtalab et al., 2019**). These enhancements could be due to existence of synergistic relationship between KS and AM (**Hajiboland et al., 2018**).

#### Fruits nutrients quality

The main effects of M inoculation and KS foliar applications as well as their interaction on squash fruits nutrients quality were illustrated in Table (6). The main effect of M inoculation was found to be positive significant for all fruit nutrients quality parameters, e.g. N, P, K, Ca, Cu, Zn and Fe, in both seasons of study. also, fruits of the M inoculated squash plants were provided more nutrient contents comparing with the fruits of the uninoculated squash plants. the average increments were estimated by 61.40, 45.62, 80.61, 28.68, 26.70, 8.54 and 12.55% for each of N, P, K, Ca, Cu, Zn and Fe, respectively for both seasons.

In case of the effect of KS foliar treatments, the KS<sub>2</sub> application was the most effective in increasing the fruit content of N, P, K, Ca, Cu and Zn, in both seasons of study. However, the effect between KS<sub>2</sub> and KS<sub>1</sub> did not differ significantly in Fe values, in both seasons. Percentage of increasing in nutrients concentrations in squash fruits due to KS<sub>2</sub> application when compared with KS<sub>0</sub> were estimated by 27.99, 25.25, 57.23, 16.02, 14.04, 5.63 and 7.74 for each of N, P, K, Ca, Cu, Zn and Fe as an average of both seasons of study, respectively.

**Table 5. Effect of mycorrhiza inoculation, potassium silicate foliar applications and their interaction on yield and its components, in 2019 and 2020 seasons**

Treatments	2019*			2020*		
	M <sub>-</sub>	M <sub>+</sub>	Mean	M <sub>-</sub>	M <sub>+</sub>	Mean
Fruits No. plant <sup>-1</sup>						
KS <sub>0</sub>	8.78 d	11.22 b	<b>10 C</b>	8.22 d	12.22 ab	<b>10.22 C</b>
KS <sub>1</sub>	9.89 c	12.22 a	<b>11.06 B</b>	9.89 c	12.44 a	<b>11.17 B</b>
KS <sub>2</sub>	10.78 bc	12.78 a	<b>11.78 A</b>	11.21 b	12.78 a	<b>12.00 A</b>
<b>Mean</b>	<b>9.81 B</b>	<b>12.07 A</b>		<b>9.78 B</b>	<b>12.48 A</b>	
Average fruit weight (gm)						
KS <sub>0</sub>	149.35 d	166.22 b	<b>157.79 C</b>	142.81 d	170.44 b	<b>156.63 C</b>
KS <sub>1</sub>	157.59 c	172.42 a	<b>165.01 B</b>	155.77 c	174.43 ab	<b>165.10 B</b>
KS <sub>2</sub>	163.44 b	177.70 a	<b>170.57 A</b>	160.74 c	178.83 a	<b>169.78 A</b>
<b>Mean</b>	<b>156.80 B</b>	<b>172.11 A</b>		<b>153.11 B</b>	<b>174.57 A</b>	
Fruit yield plant <sup>-1</sup> (kg)						
KS <sub>0</sub>	1.26 e	1.77 bc	<b>1.51 C</b>	1.21 e	1.74 bc	<b>1.49 C</b>
KS <sub>1</sub>	1.49 d	1.88 ab	<b>1.69 B</b>	1.52 d	1.91 b	<b>1.72 B</b>
KS <sub>2</sub>	1.62 cd	2.10 a	<b>1.86 A</b>	1.62 cd	2.06 a	<b>1.85 A</b>
<b>Mean</b>	<b>1.46 B</b>	<b>1.92 A</b>		<b>1.45 B</b>	<b>1.90 A</b>	
Fruit yield faddan <sup>-1</sup> (ton)						
KS <sub>0</sub>	10.09 e	14.49 bc	<b>12.29 C</b>	9.68 e	13.97 bc	<b>11.83 C</b>
KS <sub>1</sub>	12.11 d	15.57 ab	<b>13.84 B</b>	11.74 d	15.27 b	<b>13.51 B</b>
KS <sub>2</sub>	13.21 cd	17.33 a	<b>15.27 A</b>	12.84 cd	17.42 a	<b>15.13 A</b>
<b>Mean</b>	<b>11.80 B</b>	<b>15.80 A</b>		<b>11.42 B</b>	<b>15.56 A</b>	

\* Mean values with the same alphabetical letter (s) did not differ significantly at  $p < 0.05$  confidence level according to LSD test.

The interaction effect of each of M with KS was found to be significant, in both seasons of study. The foliar application of KS<sub>2</sub> under M inoculation (M<sub>+</sub>+KS<sub>2</sub>) was found to be the most pronounced than KS<sub>1</sub> or KS<sub>0</sub> in both seasons of study in the case of squash fruits content of N, P, K and Cu, in both seasons. On the other hand, the KS foliar application levels under M inoculation did not affect the nutrients content of fruits of Ca, Zn and Fe, in both seasons of study.

However, the M<sub>+</sub>+KS<sub>2</sub> treatment was still giving the highest mean values of fruits nutrients content comparing with the corresponding mean values of the squash plants grown in saline conditions without receiving any ameliorative treatment. The increasing of squash fruits nutrients content due to M<sub>+</sub>+KS<sub>2</sub> treatment were 105.69, 83.05, 205.28, 59.09, 46.71 15.49 and 22.98 for N, P, K, Ca, Cu, Zn and Fe as an average of both seasons of study, respectively.

Many of other research papers demonstrated the vital role of mycorrhiza inoculation in increasing the nutrient uptake (Colla *et al.*, 2008; Goussous and Mohammad, 2009; Bhattacharjee and Sharma, 2015; Amballa and Bhumi, 2017).

The foliar application with KS was found, in many research articles, to be appropriate treatments for enhancing the nutrient quality of squash fruits (Abd-Elaziz *et al.*, 2019; Rashad *et al.*, 2019).

Moreover, the interaction between M and KS was found to be the most superior than the single treatments, which pointed to some kind of synergistic effect between M and KS as emphasized through the research papers of Hajiboland *et al.* (2018) and Moradtalab *et al.* (2019)

According to the results of this investigation, it could be concluded that growing of squash plants c.v. "Aziad" under saline irrigation water (3.4-3.6 dSm<sup>-1</sup>) resulted in decreasing all growth, yield and fruit nutrients quality characters as shown in control. However, the application of mycorrhiza inoculum in squash seeds a combined with foliar application of potassium silicate at 4 mL<sup>-1</sup> every 10 days after transplanting in the open field could be an effective procedure for increasing the vegetative growth of squash plants grown under similar unfavorable conditions of increasing irrigation water salinity.

**Table 6.** Effect of mycorrhiza inoculation, potassium silicate foliar applications and their interaction on fruit content of N, P, K, Ca, Cu, Zn and Fe, in 2019 and 2020 seasons

Treatments	2019*			2020*		
	M <sub>-</sub>	M <sub>+</sub>	Mean	M <sub>-</sub>	M <sub>+</sub>	Mean
	N%					
KS <sub>0</sub>	0.903 e	1.542 c	<b>1.222 C</b>	.843 d	1.220 c	<b>1.032 C</b>
KS <sub>1</sub>	1.059 d	1.753 b	<b>1.406 B</b>	.941 d	1.460 b	<b>1.201 B</b>
KS <sub>2</sub>	1.189 d	1.929 a	<b>1.559 A</b>	.983 d	1.667 a	<b>1.325 A</b>
Mean	<b>1.050 B</b>	<b>1.741 A</b>		<b>0.923 B</b>	<b>1.449 A</b>	
	P%					
KS <sub>0</sub>	0.361 e	0.550 c	<b>0.457 C</b>	0.364 d	0.480 c	<b>0.422 C</b>
KS <sub>1</sub>	0.398 d	0.956 b	<b>0.497 B</b>	0.416 cd	0.557 b	<b>0.486 B</b>
KS <sub>2</sub>	0.415 d	0.671 a	<b>0.534 A</b>	0.471 c	0.656 a	<b>0.564 A</b>
Mean	<b>0.391 B</b>	<b>0.609 A</b>		<b>0.417 B</b>	<b>0.565 A</b>	
	K%					
KS <sub>0</sub>	1.626 f	3.387 c	<b>2.506 C</b>	1.424 f	3.197 c	<b>2.311 C</b>
KS <sub>1</sub>	2.412 e	4.144 b	<b>3.278 B</b>	2.143 e	3.941 b	<b>3.042 B</b>
KS <sub>2</sub>	3.039 d	4.746 a	<b>3.983 A</b>	2.651 d	4.538 a	<b>3.594 A</b>
Mean	<b>2.359 B</b>	<b>4.092 A</b>		<b>2.073 B</b>	<b>3.892 A</b>	
	Ca%					
KS <sub>0</sub>	0.225 d	0.374 a	<b>0.300 C</b>	0.249 d	0.351 a	<b>0.299 C</b>
KS <sub>1</sub>	0.271 c	0.374 a	<b>0.323 B</b>	0.278 c	0.351 a	<b>0.315 B</b>
KS <sub>2</sub>	0.336 b	0.387 a	<b>0.362 A</b>	0.301 b	0.364 a	<b>0.333 A</b>
Mean	<b>0.278 B</b>	<b>0.378 A</b>		<b>0.276 B</b>	<b>0.355 A</b>	
	Cu (ppm)					
KS <sub>0</sub>	4.072 d	5.722 b	<b>4.897 C</b>	4.111 d	4.728 bc	<b>4.420 C</b>
KS <sub>1</sub>	4.416 cd	5.971 b	<b>5.193 B</b>	4.455 c	4.913 b	<b>4.684 B</b>
KS <sub>2</sub>	4.600 c	6.788 a	<b>5.694 A</b>	4.673 bc	5.209 a	<b>4.941 A</b>
Mean	<b>4.362 B</b>	<b>6.160 A</b>		<b>4.413 B</b>	<b>4.950 A</b>	
	Zn (ppm)					
KS <sub>0</sub>	42.016 d	47.303 ab	<b>44.660 C</b>	41.556 c	45.948 ab	<b>43.752 b</b>
KS <sub>1</sub>	44.783 c	48.712 a	<b>46.748 B</b>	44.108 b	46.694 a	<b>45.401 A</b>
KS <sub>2</sub>	45.683 bc	49.144 a	<b>47.41 A</b>	44.579 b	47.375 a	<b>45.978 A</b>
Mean	<b>44.161 B</b>	<b>48.387 A</b>		<b>43.414 B</b>	<b>46.673 A</b>	
	Fe (ppm)					
KS <sub>0</sub>	43.521 c	50.894 a	<b>47.208 B</b>	42.277 d	50.744 ab	<b>46.511 B</b>
KS <sub>1</sub>	47.350 b	52.458 a	<b>49.904 A</b>	46.708 c	51.365 a	<b>49.036 A</b>
KS <sub>2</sub>	47.838 b	53.014 a	<b>50.426 A</b>	48.594 bc	52.481 a	<b>50.537 A</b>
Mean	<b>46.237 B</b>	<b>52.122 A</b>		<b>45.860 B</b>	<b>51.530 A</b>	

\* Mean values with the same alphabetical letter (s) did not differ significantly at  $p < 0.05$  confidence level according to LSD test.

## REFERENCES

Abd-Elaziz, S. A.; Alkharpotly, A. A.; Yousry, M. M. and Abido, A. I. A. (2019). Effect of foliar application with salicylic acid and potassium silicate on squash plants (*Cucurbita pepo* L.) Yield and quality. Fayoum J. Agric. Res. and Development, 33 (1), 1-30.

Abdel Latef; Arafat, A. H. and Chaoping, H. (2011). Effect of arbuscular mycorrhizal fungi on growth, mineral nutrition, antioxidant enzymes activity and fruit yield of tomato grown under salinity stress. Scientia Horticulturae, 127 (3), 228-233.

Almeida, D. M.; Oliveira, M. M. and Saibo, N. J. M. (2017). Regulation of Na<sup>+</sup> and K<sup>+</sup> homeostasis

in plants: towards improved salt stress tolerance in crop plants. Genetics and Molecular Biology, 40 (1), 326-345.

Amballa, H. and Bhumi, N. R. (2017). Significance of arbuscular mycorrhizal fungi and rhizosphere microflora in plant growth and nutrition. In Plant-Microbe Interaction: An Approach to Sustainable Agriculture (pp. 417-452).

Amirossadat, Z.; Ghehsareh, A. M. and Mojiri, A. (2012). Impact of silicon on decreasing of salinity stress in greenhouse cucumber (*Cucumis sativus* L.) in soilless culture. Journal of Biological and Environmental Sciences, 6 (17), 171-174.

Ayers, R. S. and Westcot, D. W. (1985). Water quality for agriculture (Vol. 29). Food and Agriculture Organization of the United Nations

Rome (FAO).

**Azam Eidi, S. E. and Najafabadi, G. B. K. (2014).** Effects of silicon application and mycorrhizal inoculation on growth and yield of tomato plants (*Lycopersicon esculentum* mill.) In reduced irrigation condition. The 1<sup>st</sup> International Conference on New Ideas in Argiculture, Islamic Azad University Khorasgan Branch, Isfahan, Iran, 1.

**Azia, F., and Stewart, K. A. (2001).** Relationships between extractable chlorophyll and SPAD values in muskmelon leaves. *Journal of Plant Nutrition*, 24 (6), 961-966.

**Begum, N.; Qin, C.; Ahanger, M. A.; Raza, S.; Khan, M. I.; Ahmed, N.; Ashraf, M. and Zhang, L. (2019).** Role of arbuscular mycorrhizal fungi in plant growth regulation: implications in abiotic stress tolerance. *Frontiers in Plant Science*, 10 (September), 1-15.

**Bhattacharjee, S. and Sharma, G. D. (2015).** Effect of arbuscular mycorrhizal fungi (AM fungi) and rhizobium on the nutrient uptake of pigeon pea plant. In *International Journal of Advanced Research*, 3 (8), 833-836.

**Cantrell, I. C. and Linderman, R. G. (2001).** Preinoculation of lettuce and onion with VA mycorrhizal fungi reduces deleterious effects of soil salinity. *Plant and Soil*, 233 (2), 269-281.

**Cavagnaro, T. R.; Bender, S. F.; Asghari, H. R. and van der Heijden, M. G. A. (2015).** The role of arbuscular mycorrhizas in reducing soil nutrient loss. *Trends in Plant Science*, 20, 283-290.

**Chhabra, R. (2017).** Soil salinity and water quality. Routledge.

**Colla, G.; Roupael, Y.; Cardarelli, M.; Tullio, M.; Rivera, C. M. and Rea, E. (2008).** Alleviation of salt stress by arbuscular mycorrhizal in zucchini plants grown at low and high phosphorus concentration. *Biology and Fertility of Soils*, 44 (3), 501-509.

**COSTAT (2005).** Version 6.4, Copyright (c), CoHort Software, 798, and Lighthouse Ave. PMB 320, Monterey, CA, 93940, USA.

**Decker, D. S. (1988).** Origin(s), evolution, and systematics of *Cucurbita pepo* (Cucurbitaceae). *Economic Botany*, 42 (1), 4-15.

**Dewis, J. and Freitas, F. (1970).** Physical and chemical methods of soil and water analysis. *FAO Soils Bulletin*, (10).

**Fekry, W. (2016).** Improving squash (*Cucurbita pepo* L.) Plant growth, sex expression and yield by foliar application of potassium and ethephon under high summer temperature conditions. In *J. Product. and Dev.*, 21.

**Frew, A.; Powell, J. R.; Allsopp, P. G.; Sallam, N. and Johnson, S. N. (2017).** Arbuscular mycorrhizal

fungi promote silicon accumulation in plant roots, reducing the impacts of root herbivory. *Plant and Soil*, 419 (1-2), 423-433.

**Garg, N. and Bhandari, P. (2016).** Silicon nutrition and mycorrhizal inoculations improve growth, nutrient status, K<sup>+</sup>/Na<sup>+</sup> ratio and yield of *Cicer arietinum* L. genotypes under salinity stress. *Plant Growth Regulation*, 78 (3), 371-387.

**Garg, N. and Chandel, S. (2010).** Arbuscular mycorrhizal networks: Process and functions. *Sustainable Agriculture*, 30 (3), 581-599.

**Garg, N. and Singh, S. (2018).** Arbuscular mycorrhiza rhizophagus irregularis and silicon modulate growth, proline biosynthesis and yield in *Cajanus cajan* L. Millsp.(pigeonpea) genotypes under cadmium and zinc stress. *Journal of Plant Growth Regulation*, 37 (1), 46-63.

**Goussous, S. J. and Mohammad, M. J. (2009).** Comparative effect of two arbuscular mycorrhizae and N and P fertilizers on growth and nutrient uptake of onions. *International Journal of Agriculture and Biology*, 11(4), 463-467.

**Haghighi, M. and Barzegar, M. R. (2017).** Effect of amino acid and mycorrhiza inoculation on sweet pepper growth under greenhouse conditions. *Iran Agricultural Research*, 36 (2), 47-54.

**Hajiboland, R.; Moradtalab, N.; Aliasghar zad, N.; Eshaghi, Z. and Feizy, J. (2018).** Silicon influences growth and mycorrhizal responsiveness in strawberry plants. *Physiology and Molecular Biology of Plants*, 24 (6), 1103-1115.

**Hammer, E. C.; Nasr, H.; Pallon, J.; Olsson, P. A. and Wallander, H. (2011).** Elemental composition of arbuscular mycorrhizal fungi at high salinity. *Mycorrhiza*, 21 (2), 117-129.

**Hussein, W. A. and Muhammed, M. M. (2017).** The Response of white eggplant plants to foliar application with boron and potassium silicate. *Assiut Journal of Agricultural Sciences*, 48 (1-1), 394-401.

**Kaushik, P. and Saini, D. K. (2019).** Silicon as a vegetable crops modulator- a review. *Plants*, 8 (6), 148.

**Korkmaz, A.; Karagöl, A.; Akinoğlu, G. and Korkmaz, H. (2018).** The effects of silicon on nutrient levels and yields of tomatoes under saline stress in artificial medium culture. *Journal of Plant Nutrition*, 41 (1), 123-135.

**Laane, H. M. (2018).** The effects of foliar sprays with different silicon compounds. *Plants*, 7 (2).

**Luyckx, M.; Hausman, J. F.; Lutts, S. and Guerriero, G. (2017).** Silicon and plants: current knowledge and technological perspectives. *Frontiers in Plant Science*, 8, 114.

**Machado, R.M.A. and Serralheiro, R. P. (2107).** Soil salinity: effect on vegetable crop growth.

- Management practices to prevent and mitigate soil salinization. *Horticulturae*, 3 (2), 30.
- Markwell, J.; Osterman, J. C. and Mitchell, J. L. (1995).** Calibration of the Minolta SPAD-502 leaf chlorophyll meter. *Photosynthesis Research*, 46 (3), 467-472.
- Merwad, A. R. M. A. (2018).** Response of yield and nutrients uptake of pea plants to silicate under sandy soil conditions. *Communications in Soil Science and Plant Analysis*, 49 (13), 1553-1562.
- Moradtab, N.; Hajiboland, R.; Aliasgharzad, N.; Hartmann, T. E. and Neumann, G. (2019).** Silicon and the association with an Arbuscular-Mycorrhizal fungus (*Rhizophagus clarus*) mitigate the adverse effects of drought stress on strawberry. *Agronomy*, 9 (1), 41.
- Plaut, Z.; Edelstein, M. and Ben-Hur, M. (2013).** Overcoming salinity barriers to crop production using traditional methods. *Critical Reviews in Plant Sciences*, 32 (4), 250-291.
- Quosy, D. E. D. E. (2019).** Mega Agricultural Projects in Egypt. In *Handbook of Environmental Chemistry*, 74, 593-604.
- Radwan, T. M. (2019).** Monitoring agricultural expansion in a newly reclaimed area in the western Nile Delta of Egypt using landsat imageries. *Agriculture*, 9 (7), 137.
- Rahman, M.; Hossain, A.; Ali, M. E.; Alam, F. and Alam, K. (2019).** Arbuscular mycorrhizal colonization increases yield of mungbean (*Vigna radiata* L.) at moderate level of salinity. *Soil Science and Plant Nutrition*, 65 (6), 579-88.
- Ramadan, M., and Mansour, F. (2019).** Response of pea plants to humic acid and spraying with microelements. *Journal of Productivity and Development*, 24 (2), 319-340.
- Rashad, A.; W. Adel Fekry and H. Wahdan. (2019).** Influence of some squash cultivars and growth stimulants on flowering, yield and fruit quality at autumn - winter season under open field conditions. *Journal of Productivity and Development*, 24 (3), 433-460.
- Römheld, V. and Kirkby, E. A. (2010).** Research on potassium in agriculture: Needs and prospects. *Plant and Soil*, 335 (1), 155-180.
- Safdar, H.; Amin, A.; Shafiq, Y.; Ali, A. and Yasin, R. (2019).** A review: Impact of salinity on plant growth. *Nature and Science*, 17 (1), 34-40.
- Savvas, D.; Giotis, D.; Chatzieustratiou, E.; Bakea, M. and Patakioutas, G. (2009).** Silicon supply in soilless cultivations of zucchini alleviates stress induced by salinity and powdery mildew infections. *Environmental and Experimental Botany*, 65 (1), 11-17.
- Shafeek, M.; Helmy, Y.; El-Tohamy, W.; and El-Abagy, H. (2013).** Changes in growth, yield and fruit quality of cucumber (*Cucumis sativus* L.) in response to foliar application of calcium and potassium nitrate under plastic house. *Aensiweb. Net*, 114-118.
- Shahid, S. A.; Zaman, M. and Heng, L. (2018).** Soil salinity: Historical perspectives and a world overview of the problem. In *Guideline for salinity assessment, mitigation and adaptation using nuclear and related techniques* (pp. 43-53). Springer.
- Shehata, M. N. and Abdelgawad, K. F. (2019).** Potassium silicate and amino acids improve growth, flowering and productivity of summer squash under high temperature condition. *J. Agric. and Environ. Sci*, 19 (2), 74-86.
- Tanveer, M. and Shabala, S. (2018).** Targeting redox regulatory mechanisms for salinity stress tolerance in crops. In *Salinity Responses and Tolerance in Plants, Volume 1* (pp. 213-234). Springer.
- Wang, M.; Zheng, Q.; Shen, Q.; Guo, S.; River, Y. and Key, J. (2013).** The critical role of potassium in plant stress response. *Agricultural Ministry Key Lab of Plant Nutrition and Fertilization in Low-Middle Reaches of the. Int. J. Mol. Sci*, 14, 7370-7390.
- Wu, Q. S.; Zou, Y. N. and Abd-Allah, E. F. (2014).** Mycorrhizal association and ROS in plants. In *Oxidative damage to plants* (pp. 453-475). Elsevier.
- Wungrampha, S.; Joshi, R.; Singla-Pareek, S. L., and Pareek, A. (2018).** Photosynthesis and salinity: are these mutually exclusive? *Photosynthetica*, 56 (1), 366-381.
- Kalra, Y.P. (Ed.). (1998).** *Handbook of reference methods for plant analysis*. CRC Press, Taylor and Francis Group, Boca Raton, FL.
- Yaghubi, K.; Vafae, Y.; Ghaderi, N. and Javadi, T. (2019).** Potassium silicate improves salinity resistant and affects fruit quality in two strawberry cultivars grown under salt stress. *Communications in Soil Science and Plant Analysis*, 50 (12), 1439-1451.
- Yan, G.; Nikolic, M.; YE, M.; Xiao, Z. and Liang, Y. (2018).** Silicon acquisition and accumulation in plant and its significance for agriculture. *Journal of Integrative Agriculture*, 17 (10), 2138-2150.
- Zhu, Y. X.; Gong, H. J. and Yin, J. L. (2019).** Role of silicon in mediating salt tolerance in plants: A review. *Plants*, 8 (6), 147.