



Article

Effect of Spraying Pota Crastal Compound and Vast Trivalioum Compound on Growth and Nutritioanl Status of Superior Grapevines

Farouk, H. Abd El-Aziz¹, Mohamed A.M. Abada² and Abd El Hafeez, Sh. Hassan^{1,*}

¹Hort. Dept. Fac. of Agric. Minia Univ., Egypt.

²Viticulture Dept. Hort. Res. Instit. ARC., Giza, Egypt.

*Corresponding author: Abdelhafezshehata10@gmail.com



Future Science Association

Available online free at
www.futurejournals.org

Print ISSN: 2692-5826

Online ISSN: 2692-5834

DOI:

10.37229/fsa.fjh.2023.01.10

Received: 27 November 2022

Accepted: 30 December 2022

Published: 10 January 2023

Publisher's Note: FA stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Abstract: This study was carried out in three seasons 2019, 2020 and 2021 to examine the effect of spraying Pota crastal or Vast trivolioum compounds each at 0.1% either once, twice or thrice on growth and vine nutritional status of Superior grapevines grown under El- Minia region. Application of Pota crastal or Vast trivolioum compounds once, twice or thrice each at 0.1% was very effective in enhancing main shoot length, leaf area, wood ripening coefficient, pruning wood weight/ vine, chlorophylls a , b , total chlorophylls, total carotenoids, N, P, K, Mg (as %) Fe, Mn and Zn (as ppm) of superior grapevines over the control treatment. Using vast trivolioum compound was favourable in enhancing some characteristics of vegetative growth and vine nutritional status rather than application of Pota crastal compound. The improvement in the growth characteristics and vine nutritional status was associated with an increase in the number of spraying times from one to three times. The best results with regard to characteristic growth and vine nutritional status of superior grapevines grown under El- Minia region were observed due to treating the vines three times during each season at growth start, just after berry setting and at one month later intervals with vast trivolioum compound at 0.1%.

Key words: Superior grapevines, Pota crastal compound, Vast trivolioum compound, growth, nutritional status.

INTRODUCTION

Superior grapevine cultivar is considered a prime and popular grapevine cv. successfully grown under Egypt conditions. Such cultivar ripens early sometimes in the last week of May. In addition, it early in ripening character which reduced competition. In spite of introducing numerous grapevines cvs recently to Egypt, such grapevine cvs is still one of the prime, main, popular and the most profitable grapevine cvs.

Fertilization of grapevines especially with all nutrients at balanced rate is really accompanied with improving growth and nutritional status of the vines (**Fraguas and Silva, 1998**).

Supplying various grapevines cvs with their requirements from macro and micronutrients had very beneficial effects on growth and vine

nutritional status through enhancing the biosynthesis of plant pigments proteins, carbohydrates, enzymes, fats, vitamins and hormones (Nijjar, 1985 and Tisdale *et al.*, 1985).

Amino acids, and seaweed extracts plays a definite role in protecting the plant cells from senescence and death as well as increasing the plant resistance to various disorders the biosynthesis of organic food, cell division and natural hormones and alleviating the stresses happened by environmental conditions (Karabanov, 1977; Samiullah *et al.*, 1988; Elade, 1992; Foyer and Lelandias, 1993; Sandermann *et al.*, 1998 and Khan *et al.*, 2003).

This study aimed to throw some light on the effect of compounds Pota crastal and vast trivolium as stimulant compound containing most macro and micronutrients as well as some antioxidants on growth and vine nutritional status of superior grapevines. Selecting the best number of sprays which was responsible for growth the best results regard to growth and vine nutritional.

MATERIAL AND METHODS

This study was carried out during 2019, 2020 and 2021 seasons on forty-two uniform in vigour 14-years old Superior grapevines.

The selected vines are grown in a private vineyard located at Al- Howarta village, Minia district, El-Minia Governorate – Egypt.

Where the texture of the soil is clay (Table 1) soil analysis was done according to the procedures that outlined by (Chapman and Pratt, 1987 and Black *et al.*, 1965).

The selected vines are planted at 2.0 x 3.0 meters apart (700 vines/ fed.). The chosen vine was trained by cane pruning system leaving 96 eyes/ vine (six fruiting canes x 14 eyes plus six renewal spurs x 2 eyes). Using Gable supporting method, winter pruning was carried out at the second week of December in three seasons during surface irrigation system was followed using Nile water containing 175 ppm EC.

Table 1. Analysis of the tested soil

Constituents	Values
Sand %	6.2
Silt %	19.8
Clay %	74.0
Texture	Clay
pH (1:2.5 extract)	7.98
EC (1: 2.5 extract) (dsm-1) 1cm /25°C	0.93
O.M.%	2.22
CaCO ₃ %	1.99
Total N %	0.11
Available P (ppm)	3.15
Available K (ppm)	428

Common horticultural practices such as fertilization, twice hoeings, irrigation, pinching and pest management were carried out as usual.

This study included the following seven treatments from application of Pota crastal and Vast trivolium each a 0.1% and frequencies.

1- Control (untreated vines).

- 2- Spraying Pota crastal compound at 0.1% once at growth start (1st week of Mar.)
- 3- Spraying Pota crastal compound at 0.1% twice at growth start (1st week of Mar.) and again just at berry setting (1st week of Apr.)
- 4- Spraying Pota crastal compound at 0.1% thrice at growth start (1st week of Mar.), again just at berry setting (1st week of Apr.) and one month later (1st week of May).
- 5- Spraying Vast trivolum compound at 0.1% once at growth start (1st week of Mar.)
- 6- Spraying Vast trivolum compound at 0.1% twice at growth start (1st week of Mar.) and again just at berry setting (1st week of Apr.)
- 7- Spraying Vast trivolum compound at 0.1% thrice at growth start (1st week of Mar.), again just at berry setting (1st week of Apr.) and one month later (1st week of May).

Each treatment was of three replicates, two vine per each (42 vines). Triton B as a wetting agent at 0.05% was added to all Pota crastal and Vast trivolum solutions.

Table 2. Analysis of Pota crastal

Parameters	Values
Amino acids %	5.0%
Some vitamins %	1.0%
K ₂ O %	36.0%
N %	5.0%
P %	2.0%
Some micro nutrients (Zn, Mn, B , Mo and Copalt)	2.0%

Table 3. Analysis of Vast trivolum

Parameters	Values	Parameters	Values
Amino acids %	3.0%	Mn %	1.0%
Algae seaweed extract %	4.0%	Mg%	2.5%
Total N %	7.0%	Boron %	0.25%
P %	6.0%	Cu %	0.2%
K ₂ O %	7.0%	Si %	0.2%
Fe %	1.5%	Copalt	0.01%
Zn %	1.0%		

Randomized complete block design (RCBD) was adopted for carrying out statistical analysis of this study (Rangaswamy, 1995).

During three seasons, the following measurements were recoded:

- 1- Vegetative growth characteristics namely:
 - main shoot length (cm).
 - leaf area (cm)² (Ahmed and Morsy, 1999).
 - Wood ripening coefficient (Bouard, 1966).
 - Pruning wood weight / vice (kg.).

2- Leaf chemical components namely:

- chlorophylls a, b, total chlorophylls and carotenoids content (mg/ 1.0 g F.W.) (Von – Wettstein, 1957).
- Content leaf N, P, K and Mg (as %) Zn, Fe and Mn (as ppm) (Cottenie *et al.*, 1982).

Statistical analysis was done treatment means were compared using New LSD at 5% (Mead *et al.*, 1993).

RESULTS AND DISCUSSION

1-Vegetative growth characteristics

It is clear from that data in Table (4) that applications of compounds Pota crastal or Vast trivolum each at 0.1% once, twice or thrice significantly was accompanied with stimulating the main shoot length, leaf area, wood ripening coefficient and pruning wood weight per vine comparing with non- application. The stimulation was associated with increasing compounds frequencies. Application of Vast trivolum compounds was superior the application of Pota crastal compound in enhancing growth characters of the vines. The maximum values of main shoot length (125.5, 126.5 and 128.0 cm), leaf area (111.5, 112.5 and 113.0 cm²); wood ripening coefficient (0.93, 0.94 and 0.95) pruning wood weight (2.82, 2.88 and 2.95 kg) during 2019, 2020 and 2021 seasons respectively were observed on the vines that receive three sprays of Vast trivolum compound. Untreated vines had the lowest values. These results were true during three seasons.

2- Leaf chemical composition

It is evident from the obtained data in Tables (5 to 7) that the eleven leaf chemical components namely chlorophyll a, chlorophyll b , total chlorophylls , total carotenoids, N, P, K, Mg, Fe, Zn, and Mn were significantly varied among the six compounds Pota crastal or Vast trivolum treatments. They were significantly enhanced with using compounds Pota crastal or vast trivolum each at 0.1% , relative to the control treatment.

The stimulation was associated with increasing compounds frequencies. Application of Vast trivolum was Superior the application of Pota crastal in enhancing on these leaf chemical components. The maximum values of chlorophyll a (5.3, 5.4 and 5.5 mg/ 1.0 g F.W.), chlorophyll b (2.8, 2.9 and 3.0 mg/ 1.0 g F.W.); (total chlorophylls (8.1, 8.3, 8.5 mg/ 1.0 g F.W.)); total carotenoids (2.2, 2.3 and 2.4 mg/ 1.0 g F.W.)), N (1.93, 1.96 and 1.99%); P (0.38, 0.39 and 0.39 %) ; K (1.31 , 1.32 and 1.33%) ; Mg (0.91, 0.92, 0.94%) , Zn (67.8, 68.0, 68.3 ppm) ; Fe (72.8, 73.5, 74.0 ppm) and Mn (73.2, 74.0 and 75.3 ppm) during three seasons, respectively were observed on the vines that received three sprays of compound Vast trivolum at 0.1%. The lowest values were recorded on untreated vines. These results were true during three seasons.

Table 4. Effect of spraying pota crastal and vast trivolum on some vegetative growth characteristics of Superior grapevines during 2019, 2020 and 2021 seasons

Treatments	Main shoot length (cm)			Leaf area (cm) ²			Wood ripening coefficient			Pruning wood weight / vine (kg.)		
	2019	2020	2021	2019	2020	2021	2019	2020	2021	2019	2020	2021
Control (untreated)	112.0	113.5	115.0	96.5	98.0	98.5	0.67	0.67	0.68	1.94	1.95	1.95
Spraying pota crastal compound at 0.1% once	115.5	117.0	118.0	99.5	100.0	101.0	0.71	0.72	0.73	2.25	2.30	2.35
Spraying pota crastal compound at 0.1% twice	119.0	120.5	121.0	104.5	105.0	106.0	0.77	0.78	0.79	2.40	2.45	2.48
Spraying pota crastal compound at 0.1% thrice	121.0	122.0	123.0	106.0	107.0	108.0	0.80	0.81	0.82	2.48	2.52	2.60
Spraying vast trivolum at 0.1% once	119.5	121.0	122.0	105.0	106.0	107.0	0.79	0.80	0.81	2.45	2.50	2.55
Spraying vast trivolum at 0.1% twice	124.0	125.5	126.0	109.5	110.0	111.0	0.91	0.92	0.93	2.75	2.80	2.85
Spraying vast trivolum at 0.1% thrice	125.5	126.5	128.0	111.5	112.5	113.0	0.93	0.94	0.95	2.82	2.88	2.95
NewLSD at5%	1.3	1.6	1.8	1.1	1.2	1.3	0.03	0.04	0.05	0.09	0.11	0.11

Table 5. Effect of spraying Pota crastal and vast trivolum on photosynthetic pigments in the leaves (mg/ g F.W.) of Superior grapevines during 2019, 2020 and 2021 seasons

Treatments	Chlorophyll a (mg/ g. F.W.)			Chlorophyll b (mg/ g. F.W.)			Total chlorophylls (mg/ g. F.W.)			Carotenoids (mg/ g. F.W.)		
	2019	2020	2021	2019	2020	2021	2019	2020	2021	2019	2020	2021
Control (untreated vines)	3.5	3.7	3.8	1.3	1.3	1.4	4.8	5.0	5.2	1.1	1.1	1.1
Spraying pota crastal compound at 0.1% once	4.2	4.3	4.4	1.7	1.8	1.8	5.9	6.1	6.2	1.4	1.4	1.5
Spraying pota crastal compound at 0.1% twice	4.6	4.7	4.8	2.0	2.2	2.3	6.6	6.9	7.1	1.7	1.7	1.8
Spraying pota crastal at 0.1% thrice	4.7	4.8	5.0	2.2	2.4	2.5	6.9	7.2	7.5	1.9	1.9	2.0
Spraying vast trivolum at 0.1% once	4.5	4.6	4.7	1.9	2.1	2.3	6.4	6.7	7.0	1.8	1.8	1.8
Spraying vast trivolum at 0.1% twice	5.1	5.2	5.4	2.6	2.6	2.7	7.7	7.8	8.1	2.1	2.2	2.3
Spraying vast trivolum at 0.1% thrice	5.3	5.4	5.5	2.8	2.9	3.0	8.1	8.3	8.5	2.2	2.3	2.4
New LSD at 5%	0.3	0.4	0.5	0.2	0.2	0.3	0.5	0.6	0.7	0.1	0.2	0.2

Table 6. Effect of spraying pota crastal and vast trivolum on percentages of N, p, K and Mg in the leaves of Superior grapevines during 2019, 2020 and 2021 seasons

Treatments	Leaf N %			Leaf P %			Leaf K %			Leaf Mg %		
	2019	2020	2021	2019	2020	2021	2019	2020	2021	2019	2020	2021
Control (untreated vines)	1.68	1.69	1.69	0.13	0.13	0.14	1.09	1.11	1.11	0.58	0.59	0.60
Spraying pota crastal compound at 0.1% once	1.72	1.73	1.74	0.21	0.22	0.23	1.17	1.18	1.1	0.66	0.67	0.68
Spraying pota crastal compound at 0.1% twice	1.77	1.78	1.79	0.28	0.29	0.30	1.21	1.22	1.23	0.75	0.76	0.77
Spraying pota crastal compound at 0.1% thrice	1.79	1.80	1.81	0.30	0.31	0.32	1.23	1.24	1.25	0.78	0.79	0.80
Spraying vast trivolum compound at 0.1% once	1.77	1.78	1.79	0.29	0.30	0.31	1.20	1.21	1.22	0.76	0.77	0.79
Spraying vast trivolum compound at 0.1% twice	1.89	1.91	1.93	0.36	0.37	0.38	1.28	1.29	1.30	0.86	0.87	0.89
Spraying vast trivolum compound at 0.1% thrice	1.93	1.96	1.99	0.38	0.39	0.39	1.31	1.32	1.33	0.91	0.92	0.94
New L.S.D. at 5%	0.04	0.05	0.06	0.002	0.002	0.003	0.02	0.02	0.03	0.03	0.03	0.03

Table 7. Effect of spraying pota crastal and vast trivolum on Fe, Mn and Zn (as ppm) in the leaves of Superior grapevines during 2019, 2020 and 2021 seasons

Treatments	Leaf Fe (ppm)			Leaf Mn (ppm)			Leaf Zn (ppm)		
	2019	2020	2021	2019	2020	2021	2019	2020	2021
Control (untreated vines)	53.5	54.0	55.0	55.0	55.8	56.0	48.5	49.0	49.0
Spraying pota crastal compound at 0.1% once	60.0	61.5	62.0	60.8	62.0	62.5	56.2	57.0	57.5
Spraying pota crastal compound at 0.1% twice	65.5	66.5	67.0	66.2	66.9	67.3	61.0	61.5	61.8
Spraying pota crastal compound at 0.1% thrice	68.0	68.5	69.0	68.5	69.0	71.0	62.5	62.8	63.0
Spraying vast trivolum compound at 0.1% once	66.0	66.8	67.5	66.4	67.1	67.8	61.5	61.8	62.0
Spraying vast trivolum compound at 0.1% twice	70.5	71.0	72.5	71.2	72.0	73.0	66.5	66.9	67.2
Spraying vast trivolum compound at 0.1% thrice	72.8	73.5	74.0	73.2	74.0	75.3	67.8	68.0	68.3
New L.S.D. at 5%	1.8	1.9	1.9	1.9	1.9	2.0	1.6	1.7	1.7

Discussion

The previous positive of Amino acids, seaweed extracts and Macro and micronutrients on growth, nutritional status of the vines and yield as well as physical and chemical characteristics of berries in different grapevines cvs might be attributed to antioxidative aspects, of these amino acids which play an important role in plant defense against oxidative stress induce by unfavourable conditions. Also, they are responsible for stimulating the biosynthesis of proteins, cytokinins natural hormones like IAA, ethylene and DNA, RNA, cell division, organic food and the plant pigments (**Vianello and Meric, 1991; Elade, 1992 and Orth *et al.*, 1993**).

These beneficial effects surely reflected on producing healthy vines. The present positive effects of amino acids on growth, nutritional status of the vines of superior grapevines are in agreement with those obtained by (Amin, 2007) on Red Roomy grapevines (Sayed – Heba, 2010) on Thompson seedless grapevines (Abd El-aal, 2012) on Thompson seedless grapevines (Mohamed, 2014) and (Akl *et al.*, 2014) on superior grapevines.

The beneficial effects of N, P, K, Mg, Fe, Mn, Zn, B and Co. on growth aspects, nutritional status of the vines, yield and berries quality might be ascribed to their essential roles in building amino acids, proteins, natural hormones, vitamins, organic acids, various enzymes, organic foods and plant cells. Their positive action on enhancing cell division and the tolerance of the vines to all unfavourable stress could add another explanation (Nijjar, 1985; Mengel and Kirkby, 1987, and Mengel *et al.*, 2001).

The results with regarding the effect of macro and micro nutrients are concordance with Flame seedless grapevines those obtained by Abd El- Wahab (2010) on Superior grapevines Ahmed *et al.*, (2011) on Thompson seedless grapevines, Akl *et al.*, (2014) on Superior grapevines Farahat (2017) on early sweet grapevines and Ebrahiem (2017) on Flame seedless grapevines.

The positive action of seaweed extracts applications on growth aspects, nutritional status of the vines and yield as well as berries of quality of Superior grapevines was attributed to the following reasons. It has higher content of some macro and micro nutrients as well as vitamins B₁, B₂, B₆, B₁₂ and natural hormones such as IAA, GA₃ and cytokinins and some, antioxidant (Kannaiyan, 2002 and Irizar- Garza *et al.*, 2003). These results concerning the effect of seaweed extract on promoting growth and fruiting of Superior grapevines are accordance with those obtained by Seleem – Basma and Abd El- Hameed (2008) and Gad El- Kareem and Abd El- Rahman (2013) on Ruby seedless grapevines, Farag (2006); Abd El- Hameed *et al.*, 2010; El- Saman, 2010 and Aly- Samar, 2015 on Flame seedless grapevines and Tony, (2016) on Superior grapevines.

Conclusion

Carrying out three sprays of compounds Vast trivolum at 0.1% gave the best results with regard to berry setting, yield and berries quality of Superior grapevines grown under El Mina- region.

REFERENCES

- Abd El- Hameed, H.M.; Abada, M.A.M. and Seleem- Basma, M. (2010). Reducing in organic N fertilizer partially by using yeast, seaweed and farmyard manure extract in Flame seedless grapevines. Minia 2nd Conf. Agric. Environ Sci., pp. 81-89.
- Abd El- Wahab, M.H. (2010). Relation of fruiting in superior grapevines with spraying sulphur, magnesium, zinc and boron, M. sc. Thesis, Fac. of Agric. Minia Univ. Egypt.
- Abd El-aal, E.E.H.A. (2012). The synergistic effects of using some nutrients as well as antioxidant substances on growth, nutritional status and productivity of Thompson seedless grapevines grown under Sohag region. Ph. D. Thesis Fac. of Agric. Sohag Univ., Egypt
- Ahmed, F. F.; Ibrahiem- Asmaa, A.; Mansour, A. E. M.; Shaaban, E. A. and El-Shamaa, M. S. (2011). Response of Thompson seedless grapevines to application of some amino acids enriched with nutrients as well as organic and biofertilization. Res. J. of Agric. and Biological Sci., 7 (2): 282-286.
- Ahmed, F.F. and Morsy, M.H. (1999). A new method for measuring leaf area in different fruit species. Minai J. Agric. Res., Dec., 19: 97-105.
- Akl, A.M.M.A., Abdelaziz, F.H., El- Sayed M.A. and Mohamed, T.M.M. (2014). Response of superior grapevines to spraying salicylic and boric acids. World Rural Observations, 6(4).
- Aly- Samar, S.H. (2015). Influence of reducing mineral nitrogen fertilizer partially by using plant compost enriched with *Spirulina platensis* algae fruiting of Flame seedless grapevines. M. Sc.

Thesis Fac. of Agric. Minia, Univ. Egypt.

Amin, M. M. A. (2007). Response of Red Roomy grapevines to application of amino acids and some micronutrients. M.Sc. Thesis Fac. of Agric. Minia Univ. Egypt.

Black, G. A.; Evans, D. D.; Ersminger, L. E.; White, J. L. and Dark, F. E. (1965). Methods of Soil Analysis. Amer. Soc. Agron. Inc. Bull. Medison, Wisconsin, U.S.A. pp. 891- 1400.

Bouard, J. (1966). Recherches physiologiques sur la vigne et en particulier sur laotment des serments. Thesis Sci. Nat. Bardeux, France p. 34.

Chapman, H.D. and Pratt, P.E. (1987). Methods of Analysis for Soil, Plant and Water. Univ. California, Div. Agric. Sci., 1, 150.

Cottenie, A.; Verlo, M.; Belghe, M. and Camerlynck, R. (1982). Chemical Analysis of Plant and Soil. Laboratory of Analytical and Agro- Chemistry. State Uni., Ghent, Belgium.

Ebrahiem, M.M.M. (2017). Promoting the yield quantitatively and quantitatively of Flame seedless grapevines by using amino acids Enriched with different nutrients. M. sc. Thesis Fac. of Agric. Minia Univ. Egypt.

El- Saman, A.Y.E (2010). Response of Flame seedless grapevines growing under El- Mataana conditions to spraying seaweed extract M. Sc. Thesis Fac. of Agic. Minia Univ., Egypt.

Elade, Y. (1992): The use of antioxidants to control gray mould (*Botrytic cinerg*) and white mould (*Scletotinia sclerotiorum*) in various crops. Plant Pathol., 141:417.

Farag, S.G. (2006). Minimizing mineral fertilizers in grapevines Farms to reduce the chemical residuals. M. sc. Thesis Institute of Enviro Stud. and Res. Ain Shams Univ. Egypt.

Farahat, I.A.M. (2017). Studies on pruning and fertilization of early and sweet grapevines growing under Minia Region condition. Ph.D. Thesis Fac. of Agric. Minia Univ., Egypt.

Foyer, C. H. and Lelandias, S. (1993). The role of ascorbate in regulation of photosynthesis. In Yamamoto, Y.; Smith, C. 11. (Ed), photosymthetic responses to the environment.

Fraguas, J.C. and Silvo, U.J. (1998). Nutrition of grapevines in tropical regions. in tropical regions. Inform Agropecuario, 19(194): 70-75.

Gad El- Kareem, M.R. and Abd El- Rahman, M.M.A. (2013). Response of Ruby seedless grapevines to foliar application of seaweed extract. Salicylic acid and roselle extract. Hort. Science Journal of Suez Canal Univ., 1: 294-303.

Irizar-Garza, N.U.G.; Vargas-Vazquez, P.; Garza-Garcia, D.; Tuty Couoh, C.; Rojas-Martinez, I.; Trujillo-Campos, A.; Garcia-Silva, R.; Aguirre-Montoya, D.; Martinez-Gonzalez, J.C.; Alvarado-Mendoza, S.; Grajeda-Cabrera, O.; Valero-Garza J. and Aguirre-Nledina, J.F. (2003). Respuesta de cultivosagricolas a los biofertilizantes en la region central de Mexico. Agric. Tec. Mex., 29: 213-225.

Kannaiyan, S. (2002). Biotechnology of Biofertilizers. Alpga. Sci. Inter. Ltd., P.O. Box 4067 Pang Bourne R. G8, UK. Pp. 1-375.

Karabanov, L. (1977). Effect of growth regulators and vitamins on growth and certain physiological processes in potato. Plant Growth Regul. Proc. Int. Symp., pp. 761.

Khan, H.; Ahmad, B.; Jaskani, M.J.; Ahmad, R.; Malik, A. (2003): Foliar application of mixture of amino Acids and seaweed (*Ascophylum nodosum*) extract improve growth and physico-chemical properties of grapes. Inter. J. Agric. & Biol., 14(3): 383-388.

Mead, R.; Currnow, R. N. and Harted, A. M. (1993) Statistical Methods in Agricultural and Experimental Biology. 2nd Ed. Chapman and Hall, London pp. 10- 44.

Mengel, K. and Kirkby, E.A. (1987). Principles of plant nutrition 4th E.d. International potash Institute, Pem, Switzerland pp. 10- 20.

- Mengel, K.E.; Kirkby, A.; Kosegarten, H. and Appel, T. (2001).** Principles of plant nutrition. 5th El Kluwer Academic publishers, Dordrecht, p. 1-311.
- Mohamed, W.B.M.F. (2014).** Effect of some amino acid, nutrient and salicylic acid treatments on Superior grapevine cv. M. Sc. Thesis Fac. of Agric. Assiut, Azhar Univ. Egypt.
- Nijjar, G. S. (1985).** Nutrition of Fruit Trees. Mrs Usha Raj. Kumar, for Kalyan Publishers, New Delhi India. pp. 282-302.
- Orth, A. B.; Sfarra, A.; Pell, R. J. and Tlen, M. (1993).** Assessing the involvement of free radicals in fungicide toxicity using x- tocopherol analogs pesticide. *Biochemistry and Physiology*, 47: 134 - 141.
- Rangaswamy, R. (1995).** Randomized complete Block Design. In *A text Book of Agricultural Statistics* New Age International publishers, pp. 281-309.
- Samiullah, S. A.; Ansari, M. M. and Afridi. R. K. (1988).** B- vitamins in relation to crop productivity. *Ind. Re. Life. Sci.*, pp. 80-92.
- Sandermann, H. S.; Ernst, D. and Wandlangeborels, H. C. (1998).** Ozone as abiotic elicitor of plant defense reaction trends. *Plant Sci.*, 47- 50.
- Sayed- Heba, F. I. (2010).** Effect of stimulant aminoquelant-Ca and zinc on yield and berries quality of Thompson seedless grapevines. M. Sc. Thesis Fac. of Agric., Minia Univ., Egypt.
- Seleem - Basma, M and Abd El- Harneed, H. M. (2008).** Effect of the stimulant aminoquelant- Ca on yield and berries quality of Thompson seedless grapevines. *Minia. J. of Agric. Res. & Develop.*, 28 (1):13 - 21.
- Tisdale, S.L.; Nelson, W.L. and Beaton. J.D. (1985).** *Soil Fertility and Fertilizer*. (4 edi.). Macmillan Publishing Company, New York, USA.
- Tony, M.S.S. (2016).** Partial replacement of inorganic N fertilizer in superior grapevines by using compost enriched with some microorganism. M. Sc. Thesis Fac. of Agric. Minia Univ., Egypt.
- Vianello, A. and Marei, F. (1991).** Generation of superoxide anion and hydrogen peroxide at the surface of plant cells. *J. Bioenergetics and Biomembranse*, 23(2): 409-423.
- Von- Wettstein, D. V. C. (1957).** Clatale and der Sumbmikro Skopisne Formwechsel de Plastids. *Experimental Cell Research*, 12 -427.